

LA-UR-21-20030

Approved for public release; distribution is unlimited.

Title: Heavy Flavor and jet studies for the future Electron-Ion Collider

Author(s): Li, Xuan

Intended for: The VI-th International Conference on the Initial Stages of
High-Energy Nuclear Collisions, 2021-01-10/2021-01-15 (Rehovot,
Israel)

Issued: 2021-01-14 (rev.1)

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Heavy Flavor and jet studies for the future Electron-Ion Collider

LA-UR-21-20030

Xuan Li (xuanli@lanl.gov)
Los Alamos National Laboratory

IS2021

The VIth International Conference on the
INITIAL STAGES
OF HIGH-ENERGY NUCLEAR
COLLISIONS





Outline

- What is the Electron-Ion Collider (EIC)?
- Why and how to measure heavy flavor at the EIC?
- Initial forward silicon tracker detector design and tracking performance.
- Open heavy flavor and jet studies in simulation.
- Summary and outlook.


The Electron-Ion Collider will bring new opportunities in high-energy nuclear physics

- The proposed Electron-Ion Collider (EIC) CD0 has been announced and the site is selected to be BNL.
- e-p collisions at the EIC:
 - (Polarized) p, d/³He beams at 41-275 GeV.
 - (Polarized) e beam at 5-18 GeV.
 - Instant luminosity $L_{\text{int}} \sim 10^{33-34} \text{ cm}^{-2}\text{sec}^{-1}$. A factor of ~ 1000 higher than HERA.
 - Bunch crossing rate: 1-10 ns.
- e-A collisions at the EIC:
 - Multiple nuclear species ($A=2-208$) and variable center of mass energies.
 - Instant luminosity $L_{\text{int}} \sim 10^{33-34} \text{ cm}^{-2}\text{sec}^{-1}$.
 - Bunch crossing rate: 1-10 ns.



Fundamental questions to be explored by the EIC

- The proposed EIC will (1) precisely study the nucleon/nuclei 3D structure, (2) help address the proton spin puzzle and (3) explore the nucleon/nuclei parton density extreme – gluon saturation.
- It will provide a clean environment to (4) explore how quarks and gluons form visible matter inside the vacuum/medium, which is referred to as the hadronization process.



Heavy quarks play a special role within the EIC science portfolio (I)


- Heavy quarks **c** (charm $M_c=1.3$ GeV), **b** (bottom $M_b=4.5$ GeV) are heavier than the proton. They are created in the initial collision and can probe the parton (quark or gluon) evolution processes inside the vacuum and the medium.

$$d\sigma_{\text{jet}} = f(x_B) \times H$$

Distribution of quarks and gluons in nucleons/nuclei

Accurately computable perturbative part

- The measured heavy flavor jet cross section contains information about the **initial nucleon/nuclear parton (quark or gluon) distributions**.



Heavy quarks play a special role within the EIC science portfolio (II)

- The measured heavy flavor jet/hadron cross section contains the information about both the **initial nucleon/nuclear parton distributions** and the **final state fragmentation processes**.


$$d\sigma_{\text{jet [hadron]}} = f(x_B) \times H [\times D(z_h)]$$

Distribution of quarks and gluons in nucleons/nuclei


Accurately computable perturbative part

Fragmentation function

Nuclear transport coefficient






gluon FF



- Heavy quark nuclear transport properties are predicted to be distinctly different from light quarks, giving unique discriminating power between different models.

How to measure heavy quarks in experiments?


- At the EIC, hadrons or jets which contain heavy quarks can be identified by detectors using their unique lifetime and masses.



Particle	Mass (GeV/c ²)	Average decay length
D [±]	1.869	312 micron
D ⁰	1.864	123 micron
B [±]	5.279	491 micron
B ⁰	5.280	456 micron

- Physics-driven detector performance requirements:


- Fine spatial resolution ($<100\text{ }\mu\text{m}$) for displaced vertex reconstruction.
- Fast timing resolution to suppress backgrounds from neighboring collisions.
- Low material budgets to maintain fine hit resolution.




Heavy flavor physics observables at the EIC to probe hadronization in medium

- Calculations done in the energy loss approach:
 - **Tremendous discriminating power** between models of energy loss and hadronization in matter.
 - Can **constrain nuclear opacities & transport properties to 20%**.
- Strong discriminating power provided by heavy flavor measurements to separate different nuclear effects.


EPJ Web of Conferences 235, 04002 (2020)



Heavy quark
fragmentation
modification in
 $e+A$ collisions




Light quark
fragmentation
modification in
 $e+A$ collisions



Conceptual design of the proposed Forward Silicon Tracking detector for the EIC


- GEANT4 simulation within the Fun4All framework:
 - Assumed mid-rapidity silicon vertex detector with 5 barrel layers based on the Monolithic Active Pixel Sensor (MAPS) type technology.
 - Forward-rapidity silicon tracking detector (FST) with $1.0 < \eta < 3.5$: 3 planes of MAPS silicon detector and 2 forward planes of HV-MAPS silicon detector.

LANL FST integrated inside the EIC




arXiv:2009.
02888

LANL FST tracking performance meets the open heavy flavor reconstruction requirements



Reconstructed heavy flavor hadron with the proposed FST in simulation

- The full analysis framework which includes the event generation (PYTHIA), detector response in GEANT4 simulation, beam remnant & QCD background, and hadron reconstruction algorithm have been setup.
- Mass distributions of reconstructed D-meson and B-meson family in 10 GeV electron and 100 GeV proton collisions with integrated luminosity: 10 fb^{-1} .







arXiv:2009.02888

- Central, backward and forward tracking performance.
- Primary vertex resolution: 20-35 (μm) depends on the track multiplicity.
- 95% $K/\pi/p$ separation over all the acceptance.
- Charged track clusters with a decay length (DCA) cut.

Flavor dependent nuclear modification factor projections for reconstructed hadrons

- Inclusive flavor dependent hadron nuclear modification factor R_{eA} projection in 10+100 GeV e+Au collisions.




- Good statistical uncertainties can be achieved by reconstructed heavy flavor hadrons.
- Can provide sufficient discriminator power to separate different model predictions.

Separate the kinematics: pseudorapidity dependence

- Heavy flavor produced in different pseudorapidity regions experience different initial and final state effects.

η dependent reconstructed D^0 mass distribution




E_e : 10 GeV
 E_p : 100 GeV
Int. Lumi: 10 fb^{-1}

[arXiv:2009.02888](https://arxiv.org/abs/2009.02888)

Separate the kinematics: pseudorapidity dependence

- Heavy flavor measurements especially in the forward regions at the EIC has enhanced sensitivity to the hadronization process in medium and the nuclear transport properties.




p_T dependent R_{eA} for D^0 meson

Theoretical calculations from the HF
tomography in EIC, arXiv: 2007.10994

Separate the kinematics: pseudorapidity dependence

- Heavy flavor measurements especially in the forward regions at the EIC has enhanced sensitivity to the hadronization process in medium and the nuclear transport properties.




z_h dependent R_{eA} for D^0 meson

Theoretical calculations from the HF
tomography in EIC, arXiv: 2007.10994

Separate the kinematics: pseudorapidity dependence

- Heavy flavor measurements especially in the forward regions at the EIC has enhanced sensitivity to the hadronization process in medium and the nuclear transport properties.




p_T depend R_{eA} for B^\pm meson

Theoretical calculations from the HF tomography in EIC, arXiv: 2007.10994

Heavy flavor hadron and jet studies


- More reconstructed heavy flavor products have been explored in the full simulation including vertex, tracking and PID performance.

Charm baryon reconstruction



A different approach to the hadronization process such as Λ_c/D ratio to check the impacts from recombination in vacuum/medium.

Flavor tagged jet yields




- Jet reconstruction using the anti- k_T algorithm with cone radius 0.8.
- Tag **charm-jets** with at least one charm hadron inside the jet cone.
- If no heavy flavor hadrons are found inside the jet cone, tag this jet as a light flavor jet.
- Jet yields are not corrected by the reconstruction efficiency yet.

Jet substructure for different flavor jets

- A new probe to explore the hadronization origin and process: jet angularity.

Definition: $\tau_a \equiv \tau_a^{pp} \equiv \frac{1}{p_T} \sum_{i \in J} p_T^i (\Delta \mathcal{R}_{iJ})^{2-a}$ **JHEP 1804 (2018) 110**




Initial studies in arXiv: 2007.14417



- Jet origin from a quark/gluon can be distinguished from this study.
- Shed light into how quark/gluon recombined into final hadrons with different masses.
- Impacts by nuclear medium effects will be studied in e+A collisions.

Summary and Outlook


- Nice progresses and results have been achieved in the EIC heavy flavor and jet studies with detector performances evaluated in full simulation.
- The new heavy flavor and jet program for the EIC will explore the flavor dependent parton energy loss in medium and the hadronization processes in the poorly constrained kinematic region.
- We look forward to work with more collaborators and contribute to the EIC realization.



Backup


New EIC heavy flavor and jet program at LANL

- An EIC DR (20200022DR), Oct. 2019 to Sep. 2022, is funded by the LANL LDRD office with PI: Ivan Vitev, Co-PI: Xuan Li and 15+ staff/postdocs.




- Through this EIC project at LANL, we will explore hadronization processes and their medium modifications using heavy flavor and jet probes at the EIC.

- We will carry out detector R&D for several advanced silicon sensor candidates and complete the conceptual design for a **forward silicon tracking detector** to realize the EIC heavy flavor and jet physics measurements.





D-meson kinematics

- In 10 GeV electron and 100 GeV proton collisions with integrated luminosity: 10 fb^{-1}




Reconstructed heavy flavor hadron with the proposed FST in simulation

- Heavy flavor physics studies can provide guidance on the EIC detector technology selection and conceptual design.
- Mass distributions of fully reconstructed D-meson and B-meson.
- Provide detector requirements based on physics needs.







Signal/Background VS Material budget per layer



Reconstructed D^0 mass distributions in different p_T bins


- In 10+100 GeV e+p collisions with integrated luminosity at 10 fb^{-1} .




Exotic heavy flavor states at the EIC

- New physics observables are under study.
 - Structure and formation process of new exotic hadrons, e.g. X(3872) can be explored by measuring their suppression in e+A collisions.


PRD 71, 014028 (2005)
PLB 662 424 (2008)



PLB 590 209 (2004)
PRD 77 014029 (2008)
PRD 100 0115029(R) (2019)



Relative modification of X(3872)/ $\psi(2S)$
projection at $\sqrt{s} = 63.2\text{GeV}$



Arleo et al., PRC, 61 054906 (2000)